Calculus 1 Workbook Solutions

Sketching graphs

krista king MATH

VERTICAL ASYMPTOTES

■ 1. Find any vertical asymptote(s) of the function.

$$f(x) = \frac{-x^2 + 16x - 63}{x^2 - 2x - 35}$$

Solution:

Factor the numerator and denominator as completely as possible.

$$f(x) = \frac{-(x - 7)(x - 9)}{(x - 7)(x + 5)}$$

The denominator is equal to 0 if $x = 7$ or $x = -5$, which means the function has two discontinuities. However, the function simplifies to

$$f(x) = \frac{-(x - 9)}{x + 5}$$

$$f(x) = \frac{9 - x}{x + 5}$$

Therefore, the function has a removable discontinuity at $x = 7$ and a vertical asymptote at $x = -5$, which means the domain of the function is $(-\infty, -5) \cup (-5,7) \cup (7,\infty)$.

■ 2. Find any vertical asymptote(s) of the function.

© Krista King Math

$$g(x) = \frac{x^2 - 3x - 10}{x^2 + x - 2}$$

Solution:

Factor the numerator and denominator as completely as possible.

$$g(x) = \frac{(x - 5)(x + 2)}{(x - 1)(x + 2)}$$

The denominator is equal to 0 if $x = -2$ or $x = 1$, which means the function has two discontinuities. However, the function simplifies to

$$g(x) = \frac{x - 5}{x - 1}$$

Therefore, the function has a removable discontinuity at $x = -2$ and a vertical asymptote at $x = 1$, which means the domain of the function is $(-\infty, -2) \cup (-2,1) \cup (1,\infty)$.

3. Find any vertical asymptote(s) of the function.

$$h(x) = \frac{40 - 27x - 12x^2 - x^3}{9x^2 + 63x - 72}$$

Solution:

Factor the numerator and denominator as completely as possible.

© Krista King Math

$$h(x) = \frac{-(x+8)(x+5)(x-1)}{9(x-1)(x+8)}$$

Cancel common factors from the numerator and denominator, then simplify.

$$h(x) = \frac{-(x+5)}{9}$$

$$h(x) = -\frac{x+5}{9}$$

There are no values of x that make this denominator 0, so the function has no vertical asymptotes. But it does have removable discontinuities for the factors we canceled, at $x = -8$ and $x = 1$.

© Krista King Math

HORIZONTAL ASYMPTOTES

■ 1. Find the horizontal asymptote(s) of the function.

$$f(x) = \frac{8x^4 - x^2 + 1}{4x^4 - 1}$$

Solution:

In a polynomial function, the term with the highest degree dominates the behavior of the function. So the behavior of $f(x)$ is dominated by the behavior of

$$\frac{8x^4}{4x^4}$$

Find the equation of the horizontal asymptote by taking the limit of the dominating function as $x \to \infty$.

$$\lim_{x\to\infty} \frac{8x^4}{4x^4} = \lim_{x\to\infty} \frac{8}{4} = 2$$

Therefore, the equation of the horizontal asymptote is $y = 2$.

■ 2. Find the horizontal asymptote(s) of the function.

$$g(x) = \frac{2x^2 - 5x + 12}{3x^2 - 11x - 4}$$

© Krista King Math

Solution:

In a polynomial function, the term with the highest degree dominates the behavior of the function. So the behavior of $g(x)$ is dominated by the behavior of

$$\frac{2x^2}{3x^2}$$

Find the equation of the horizontal asymptote by taking the limit of the dominating function as $x \to \infty$.

$$\lim_{x\to\infty} \frac{2x^2}{3x^2} = \lim_{x\to\infty} \frac{2}{3} = \frac{2}{3}$$

Therefore, the equation of the horizontal asymptote is $y = 2/3$.

■ 3. Find the horizontal asymptote(s) of the function.

$$h(x) = \frac{x^3 - x^2 + 6x - 1}{7x^4 - 1}$$

Solution:

In a polynomial function, the term with the highest degree dominates the behavior of the function. So the behavior of $g(x)$ is dominated by the behavior of

© Krista King Math

$$\frac{x^3}{7x^4}$$

Find the equation of the horizontal asymptote by taking the limit of the dominating function as $x \to \infty$.

$$\lim_{x\to\infty} \frac{x^3}{7x^4} = \lim_{x\to\infty} \frac{1}{7x} = 0$$

Therefore, the equation of the horizontal asymptote is $y = 0$.

© Krista King Math

SLANT ASYMPTOTES

■ 1. Find the slant asymptote of the function.

$$f(x) = \frac{3x^4 - x^3 + x^2 - 4}{x^3 - x^2 + 1}$$

Solution:

Use polynomial long division on the function to rewrite it as

$$3x + 2 + \frac{3x^2 - 3x - 6}{x^3 - x^2 + 1}$$

Then take the limit of the rational portion of the quotient as $x \to \infty$.

$$\lim_{x\to\infty} \frac{3x^2 - 3x - 6}{x^3 - x^2 + 1} = \lim_{x\to\infty} \frac{3x^2}{x^3} = \lim_{x\to\infty} \frac{3}{x} = 0$$

Therefore, the equation of the slant asymptote is $y = 3x + 2$.

■ 2. Find the slant asymptote of the function.

$$g(x) = \frac{8x^2 + 14x - 7}{4x - 1}$$

Solution:

© Krista King Math

Use polynomial long division on the function to rewrite it as

$$2x + 4 - \frac{3}{4x - 1}$$

Then take the limit of the rational portion of the quotient as $x \to \infty$.

$$\lim_{x\to\infty} -\frac{3}{4x - 1} = \lim_{x\to\infty} -\frac{3}{4x} = 0$$

Therefore, the equation of the slant asymptote is $y = 2x + 4$.

■ 3. Find the slant asymptote of the function.

$$h(x) = \frac{x^3 - 8}{x^2 - 5x + 6}$$

Solution:

Use polynomial long division on the function to rewrite it as

$$x + 5 + \frac{19x - 38}{x^2 - 5x + 6}$$

Then take the limit of the rational portion of the quotient as $x \to \infty$.

$$\lim_{x\to\infty} \frac{19x - 38}{x^2 - 5x + 6} = \lim_{x\to\infty} \frac{19x}{x^2} = \lim_{x\to\infty} \frac{19}{x} = 0$$

© Krista King Math

Therefore, the equation of the slant asymptote is $y = x + 5$.

© Krista King Math

SKETCHING GRAPHS

■ 1. Sketch the graph of the function.

$f(x) = x^3 - 4x^2 + 8.$

Solution:

Take the derivative, then set it equal to 0 to find critical points.

$$f'(x) = 3x^2 - 8x$$

$$3x^2 - 8x = 0$$

$$x(3x - 8) = 0$$

$$x = 0, \frac{8}{3}$$

Use the first derivative test to see where $f(x)$ is increasing and decreasing.

Interval	x<0	x=0	0<x<8/3	x=8/3	x>8/3
x	-2	0	1	8/3	4
f'(x)	+	0	-	0	+
Direction	Increasing	Maximum	Decreasing	Minimum	Increasing

We can see that $f(x)$

- increases on the interval $(-\infty,0)$,

© Krista King Math

- has a local maximum at $x = 0$,
- decreases on the interval $(0,8/3)$,
- has a local minimum at $x = 8/3$, and then
- increases on the interval $(8/3,\infty)$.

Evaluate the function at the extrema.

$$f(0) = (0)^3 - 4(0)^2 + 8 = 8$$

$$f\left(\frac{8}{3}\right) = \left(\frac{8}{3}\right)^3 - 4\left(\frac{8}{3}\right)^2 + 8 = -\frac{40}{27}$$

There's a local maximum at $(0,8)$ and a local minimum at $(8/3, -40/27)$. Now use the second derivative to determine concavity.

$$f''(x) = 6x - 8$$

$$6x - 8 = 0$$

$$x = \frac{4}{3}$$

Test values around the inflection point $x = 4/3$.

Interval	x<4/3	x=4/3	x>4/3
x	0	4/3	3
f''(x)	-	0	+
Concavity	Down	Inflection	Up

© Krista King Math

We can see that $f(x)$ is concave down on the interval $(-\infty,4/3)$ and concave up on the interval $(4/3,\infty)$. Because $f(4/3) = 88/27$, $f(x)$ has an inflection point at $(4/3,88/27)$. Since $f(x)$ is a polynomial function, its graph has no asymptotes.

Putting all this together, the graph is



2. Sketch the graph of the function.

$$g(x) = \frac{1}{4}x^4 - \frac{1}{3}x^3 - 3x^2 + 1$$

Solution:

Take the derivative, then set it equal to 0 to find critical points.

© Krista King Math

$$g'(x) = x^3 - x^2 - 6x$$

$$x^3 - x^2 - 6x = 0$$

$$x(x-3)(x+2) = 0$$

$$x = -2,\ 0,\ 3$$

Use the first derivative test to see where $g(x)$ is increasing and decreasing.

Interval	x<-2	x=-2	-2<x<0	x=0	0<x<3	x=3	x>3
x	-4	-2	-1	0	2	3	4
g'(x)	-	0	+	0	-	0	+
Direction	Decreasing	Minimum	Increasing	Maximum	Decreasing	Minimum	Increasing

We can see that $g(x)$

- decreases on the interval $(-\infty, -2)$,
- has a local minimum at $x = -2$,
- increases on the interval $(-2,0)$,
- has a local maximum at $x = 0$
- decreases on the interval $(0,3)$
- has a local minimum at $x = 3$, and then
- increases on the interval $(3,\infty)$.

Evaluate the function at the extrema.

© Krista King Math

$$g(-2) = \frac{1}{4}(-2)^4 - \frac{1}{3}(-2)^3 - 3(-2)^2 + 1 = -\frac{13}{3}$$

$$g(0) = \frac{1}{4}(0)^4 - \frac{1}{3}(0)^3 - 3(0)^2 + 1 = 1$$

$$g(3) = \frac{1}{4}(3)^4 - \frac{1}{3}(3)^3 - 3(3)^2 + 1 = -\frac{59}{4}$$

There's a local minimum at $(-2, -13/3)$, a local maximum at $(0,1)$, and a local minimum at $(3, -59/4)$. Now use the second derivative to determine concavity.

$$g''(x) = 3x^2 - 2x - 6$$

$$3x^2 - 2x - 6 = 0$$

$$x = \frac{2 \pm \sqrt{4 + 72}}{6} = \frac{1 \pm \sqrt{19}}{3}$$

Test values around the inflection points using their approximate values.

Interval	x<-1.12	x=-1.12	-1.12<x<1.79	x=1.79	x>1.79
x	-2	-1.2	1	1.79	4
g''(x)	+	0	-	0	+
Concavity	Up	Inflection	Down	Inflection	Up

We can see that $g(x)$ is concave up on the interval $(-\infty, (1 - \sqrt{19})/3)$, concave down on the interval $((1 - \sqrt{19})/3, (1 + \sqrt{19})/3)$, and concave up on the interval $((1 + \sqrt{19})/3, \infty)$. Because $g((1 - \sqrt{19})/3) \approx -1.9$, $g(x)$ has an inflection point at approximately $(-1.12, -1.9)$. Because $g((1 + \sqrt{19})/3) \approx -7.96$, $g(x)$ has an inflection point at approximately

© Krista King Math

$(1.79, -7.96)$. Since $g(x)$ is a polynomial function, its graph has no asymptotes.

Putting all this together, the graph is



■ 3. Sketch the graph of the function.

$$h(x) = \frac{x^2 + x - 6}{4x^2 + 16x + 12}$$

Solution:

Take the derivative, then set it equal to 0 to find critical points.

© Krista King Math

$$h'(x) = \frac{12(x^2 + 6x + 9)}{(4x^2 + 16x + 12)^2} = \frac{12(x+3)^2}{\left[4(x+1)(x+3)\right]^2} = \frac{12(x+3)^2}{16(x+1)^2(x+3)^2} = \frac{3}{4(x+1)^2}$$

There are no values for which $h'(x) = 0$, so there are no critical points. But $h'(x)$ is undefined when $x = -1$. Now use the second derivative to determine concavity.

$$h''(x) = -\frac{3}{2(x+1)^3}$$

There are no values for which $h''(x) = 0$, but $h''(x)$ is undefined when $x = -1$. Test values around the inflection point $x = -1$.

Interval	x<-1	x=-1	x>-1
x	-3	-1	3
h”(x)	+	DNE	-
Concavity	Up	Inflection	Down

We can see that $h(x)$ is concave up on the interval $(-\infty, -1)$ and concave down on the interval $(-1, \infty)$. Since $h(x)$ is a rational function, we need to look for asymptotes.

The behavior of the function is dominated by the highest degree terms in the numerator and denominator, which means the horizontal asymptote is

$$\lim_{x \to \pm\infty} \frac{x^2}{4x^2} = \lim_{x \to \pm\infty} \frac{1}{4} = \frac{1}{4}$$

The denominator of $h(x)$ is 0 when $x = -1$, so the function has a vertical asymptote there. To determine the behavior of the function near $x = -1$, find

© Krista King Math

$$\lim_{x \to -1^-} h(x) = \infty$$

$$\lim_{x \to -1^+} h(x) = -\infty$$

Lastly, the function crosses the x-axis when the numerator equals 0, which occurs at $x = 2$, and the function crosses the y-axis when $x = 0$, which gives a y-intercept of $(0, -1/2)$.

Putting all this together, the graph is



© Krista King Math

EXTREMA ON A CLOSED INTERVAL

■ 1. Find the extrema of $f(x) = x^3 - 3x^2 + 5$ over the closed interval $[-3,4]$.

Solution:

Find the critical points of the function.

$$f'(x) = 3x^2 - 6x$$

$$3x^2 - 6x = 0$$

$$3x(x - 2) = 0$$

$$x = 0,\ 2$$

Evaluate the function at the endpoints and at the critical numbers.

For $x = -3$, $(-3)^3 - 3(-3)^2 + 5 = -27 - 27 + 5 = -49$

For $x = 0$, $(0)^3 - 3(0)^2 + 5 = 0 - 0 + 5 = 5$

For $x = 2$, $(2)^3 - 3(2)^2 + 5 = 8 - 12 + 5 = 1$

For $x = 4$, $(4)^3 - 3(4)^2 + 5 = 64 - 48 + 5 = 21$

The results show that $f(x)$ has a global minimum at $(-3, -49)$, a local maximum at $(0,5)$, a local minimum at $(2,1)$, and a global maximum at $(4,21)$.

© Krista King Math

■ 2. Find the extrema of $g(x) = \sqrt[3]{2x^2 + 3}$ over the closed interval $[-1,5]$.

Solution:

Find the critical points of the function.

$$g'(x) = \frac{1}{3}(2x^2 + 3)^{-\frac{2}{3}}(4x) = \frac{4x}{3\sqrt[3]{(2x^2 + 3)^2}}$$

$$4x = 0$$

$$x = 0$$

Evaluate the function at the endpoints and at the critical numbers.

For $x = -1$, $\sqrt[3]{2(-1)^2 + 3} = \sqrt[3]{5} \approx 1.71$

For $x = 0$, $\sqrt[3]{2(0)^2 + 3} = \sqrt[3]{3} \approx 1.44$

For $x = 5$, $\sqrt[3]{2(5)^2 + 3} = \sqrt[3]{53} \approx 3.76$

The results show that $g(x)$ has a global minimum at $(0,\sqrt[3]{3})$, and a global maximum at $(5,\sqrt[3]{53})$.

■ 3. Find the extrema of $h(x) = -4x^3 + 6x^2 - 3x - 2$ over the closed interval $[-4,6]$.

© Krista King Math

Solution:

Find the critical points of the function.

$$h'(x) = -12x^2 + 12x - 3$$

$$-12x^2 + 12x - 3 = 0$$

$$-3(4x^2 - 4x + 1) = 0$$

$$-3(2x - 1)(2x - 1) = 0$$

$$x = 1/2$$

Evaluate the function at the endpoints and at the critical numbers.

For $x = -4$, $\quad -4(-4)^3 + 6(-4)^2 - 3(-4) - 2 = 362$

For $x = 1/2$, $\quad -4(1/2)^3 + 6(1/2)^2 - 3(1/2) - 2 = -5/2$

For $x = 6$, $\quad -4(6)^3 + 6(6)^2 - 3(6) - 2 = -668$

The results show that $h(x)$ has a global maximum at $(-4,362)$, a horizontal tangent line at $(1/2, -5/2)$, and a global minimum at $(6, -668)$.

© Krista King Math

SKETCHING F(X) GIVEN THE GRAPH OF F'(X), OR VICE VERSA

■ 1. Sketch a possible graph of $f(x)$ given the graph below of $f'(x)$.



Solution:

The graph of $f'(x)$ is below the x-axis on the intervals $(-\infty,0)$ and $(2,\infty)$, which means the function $f(x)$ has a negative slope and is decreasing on these intervals.

Additionally, the graph of $f'(x)$ is above the x-axis on the interval $(0,2)$, which means the function $f(x)$ has a positive slope and is increasing on this interval.

© Krista King Math

The graph of $f'(x)$ passes through the x-axis and changes sign from negative to positive at $x = 0$, which means that the graph of $f(x)$ has a minimum value at $x = 0$, and the graph of $f'(x)$ passes through the x-axis and changes sign from positive to negative at $x = 2$, which means that the graph of $f(x)$ has a maximum value at $x = 2$.

The graph of $f'(x)$ has a maximum value at $x = 1$, and its slope changes from positive to negative at that point. This means that the graph of $f(x)$ is concave up to the left of $x = 1$, has an inflection point at $x = 1$, and is concave down to the right of $x = 1$.

Putting these facts together, and based on the "assumption" that $f(x)$ contains the point $(0,0)$, this is a possible graph of $f(x)$:



© Krista King Math

2. Sketch a possible graph of $g'(x)$ given the graph below of $g(x)$.



Solution:

The graph of $g(x)$ has a positive slope on the intervals $(-\infty, -1.5)$ and $(0,1.5)$. Since $g'(x)$ is the derivative of $g(x)$, the graph of $g'(x)$ is above the x-axis on these intervals.

The graph of $g(x)$ has a negative slope on the intervals $(-1.5,0)$ and $(1.5,\infty)$. Since $g'(x)$ is the derivative of $g(x)$, the graph of $g'(x)$ is below the x-axis on these intervals.

The graph of $g(x)$ has a maximum value at $x = -1.5$ and $x = 1.5$ and its slope is 0, so the graph of $g'(x)$ passes through the x-axis and changes sign from positive to negative at $x = -1.5$ and $x = 1.5$.

© Krista King Math

The graph of $g(x)$ has a minimum value at $x = 0$, and its slope changes from negative to positive at that point. This means that the graph of $g'(x)$ passes through the x-axis at $x = 0$, and changes from negative to positive.

It appears that the graph of $g(x)$ has an inflection point at $x = -0.75$ and $x = 0.75$, so the graph of $g'(x)$ has extrema at those points. Putting these facts together, this is a possible graph of $g'(x)$:



■ 3. Sketch a possible graph of $h(x)$ given the graph below of $h'(x)$.

© Krista King Math



Solution:

The graph of $h'(x)$ is above the x-axis on the intervals $(-\infty, -3)$ and $(-3,\infty)$, which means the function $h(x)$ has positive slopes and is increasing on these intervals. Since we're only excluding the single point $x = -3$, that means the function is essentially increasing everywhere.

The graph of $h'(x)$ is on the x-axis at $x = -3$, which means the function $h(x)$ has a horizontal tangent at $x = -3$ and is increasing on both sides of this point.

The graph of $h'(x)$ has a minimum value at $x = -3$, and its slope changes from positive to negative at that point. This means that the graph of $h(x)$ is concave down to the left of $x = -3$, has an inflection point at $x = -3$, and is

© Krista King Math

concave up to the right of $x = -3$. Putting these facts together, this is a possible graph of $h(x)$:



■ 4. Sketch a possible graph of $f'(x)$ given the graph below of $f(x)$.

© Krista King Math



Solution:

The graph of $f(x)$ has a positive slope on the interval $(-4,0)$. Since $f'(x)$ is the derivative of $f(x)$, the graph of $f'(x)$ is above the x-axis on this interval.

The graph of $f(x)$ has a negative slope on the interval $(0,4)$. Since $f'(x)$ is the derivative of $f(x)$, the graph of $f'(x)$ is below the x-axis on this interval.

The graph of $f(x)$ has a maximum value at $x = 0$ and its slope is 0, so the graph of $f'(x)$ passes through the x-axis and changes sign from positive to negative at $x = 0$.

The graph of $f(x)$ has vertical tangents at $x = -4$ and $x = 4$. This means that the graph of $f'(x)$ has these limits:

$$\lim_{x \to -4^+} f'(x) = \infty$$

© Krista King Math

$$\lim_{x \to 4^-} f'(x) = -\infty$$

The graph of $f(x)$ has no inflection points so the graph of $f'(x)$ has no extrema in the interval $(-4,4)$. Putting these facts together, this is a possible graph of $f'(x)$:



■ 5. Sketch a possible graph of $f(x)$ given the graph below of $f'(x)$.

© Krista King Math



Solution:

The graph of $f'(x)$ is below the x-axis on the interval $(-\infty,0)$, which means the function $f(x)$ has a negative slope and is decreasing on this interval.

The graph of $f'(x)$ is above the x-axis on the interval $(0,\infty)$, which means the function $f(x)$ has a positive slope and is increasing on this interval.

The graph of the $f'(x)$ does not pass through the x-axis, which means that the graph of $f(x)$ does not have any extrema.

The slope of the graph of $f'(x)$ is negative on $(-\infty,0)$ and $(0,\infty)$. This means that the graph of $f(x)$ is concave down to the left and to the right of the y-axis.

© Krista King Math

The graph of $f'(x)$ has these limits:

$$\lim_{x \to 0^-} f'(x) = -\infty$$

$$\lim_{x \to 0^+} f'(x) = \infty$$

This means the graph of $f(x)$ has an asymptote on the y-axis. Putting these facts together, this is a possible graph of $f(x)$:



■ 6. Sketch a possible graph of $g'(x)$ given the graph below of $g(x)$.

© Krista King Math



Solution:

The graph of $g(x)$ has a positive slope on the intervals $(-\infty,3)$ and $(5,\infty)$. Since $g'(x)$ is the derivative of $g(x)$, the graph of $g'(x)$ is above the x-axis on these intervals.

The graph of $g(x)$ has a negative slope on the interval the $(3,5)$. Since $g'(x)$ is the derivative of $g(x)$, the graph of $g'(x)$ is below the x-axis on this interval.

The graph of $g(x)$ has a maximum value at $x = 3$ and its slope is 0, so the graph of $g'(x)$ passes through the x-axis and changes sign from positive to negative at $x = 3$.

The graph of $g(x)$ has a minimum value at $x = 5$, and its slope changes from negative to positive at that point. This means that the graph of $g'(x)$ passes through the x-axis at $x = 5$, and changes from negative to positive.

© Krista King Math

It appears that the graph of $g(x)$ has an inflection point at $x = 4$, so the graph of $g'(x)$ has extrema at $x = 4$. Putting these facts together, this is a possible graph of $g'(x)$:





WWW.KRISTAKINGMATH.COM